

Electron paramagnetic resonance in mixed crystals (BaF₂)_{1-x}(LaF₃)_x Activated by Ce³⁺ Ions

Aminov L., Abdulsabirov R., Korableva S., Kurkin I., Kurzin S., Ziganshin A., Gromov I.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The electron paramagnetic resonance (EPR) spectra of mixed crystals (BaF₂)_{1-x-y}(LaF₃)_x(CeF₃)_y ($y = 0.001 = 0.1\%$, $x = 0-0.02$) are investigated in a magnetic field $H \parallel C_4$ at a frequency of 9.5 GHz. The angular dependence of the EPR spectrum is measured for the sample with $x = 0.02$. The lines attributed to Ce³⁺ impurity centers with tetragonal symmetry and g factors ($g_{\parallel} = 0.75$, $g_{\perp} = 2.4$) close to those measured for the KY₃F₁₀: Ce³⁺ compound are separated in the complex EPR spectrum. The assumption is made that the aforementioned impurity centers are cubooctahedral clusters of the La₆F₃₇ type in which one of the La³⁺ ions is replaced by the Ce³⁺ ion. © 2005 Pleiades Publishing, Inc.

<http://dx.doi.org/10.1134/1.2014492>
